

# CME 307 / MS&E 311: Optimization

## Operators

Professor Udell

Management Science and Engineering  
Stanford

May 10, 2023

# Announcements



# Outline

## Subgradients

Subgradient properties

Subgradient method

Proximal operators

Proximal gradient method

Relations

Fixed points

Averaged operators

## Basic inequality

recall basic inequality for convex differentiable  $f$ :

$$f(y) \geq f(x) + \nabla f(x)^T(y - x)$$

- ▶ first-order approximation of  $f$  at  $x$  is global underestimator
- ▶  $(\nabla f(x), -1)$  supports **epi**  $f$  at  $(x, f(x))$

what if  $f$  is not differentiable?

## Non-differentiable functions

are these functions differentiable?

- ▶  $|t|$  for  $t \in \mathbf{R}$
- ▶  $\|x\|_1$  for  $x \in \mathbf{R}^n$
- ▶  $\|X\|_*$  for  $X \in \mathbf{R}^{n \times n}$
- ▶  $\max_i a_i^T x + b_i$  for  $x \in \mathbf{R}^n$
- ▶  $\lambda_{\max}(X)$  for  $X \in \mathbf{R}^{n \times n}$
- ▶ indicators of convex sets  $\mathcal{C}$

if not, where? can we find underestimators for them?

## Subgradient of a function

$g$  is a **subgradient** of  $f$  (not necessarily convex) at  $x$  if

$$f(y) \geq f(x) + g^T(y - x) \quad \text{for all } y$$

picture

## Subgradient of a function

$g$  is a **subgradient** of  $f$  (not necessarily convex) at  $x$  if

$$f(y) \geq f(x) + g^T(y - x) \quad \text{for all } y$$

picture

**Q:** Can a function  $f$  have  $> 1$  subgradient at a point  $x$ ?

## Subgradient of a function

$g$  is a **subgradient** of  $f$  (not necessarily convex) at  $x$  if

$$f(y) \geq f(x) + g^T(y - x) \quad \text{for all } y$$

picture

**Q:** Can a function  $f$  have  $> 1$  subgradient at a point  $x$ ?

**A:** Yes, if  $f$  is nonsmooth at  $x$



## Subgradient of a function

$g$  is a **subgradient** of  $f$  (not necessarily convex) at  $x$  if

$$f(y) \geq f(x) + g^T(y - x) \quad \text{for all } y$$

picture

**Q:** Can a function  $f$  have  $> 1$  subgradient at a point  $x$ ?

**A:** Yes, if  $f$  is nonsmooth at  $x$

**Q:** Can a function  $f$  have no subgradient at a point  $x$ ?

## Subgradient of a function

$g$  is a **subgradient** of  $f$  (not necessarily convex) at  $x$  if

$$f(y) \geq f(x) + g^T(y - x) \quad \text{for all } y$$

picture

**Q:** Can a function  $f$  have  $> 1$  subgradient at a point  $x$ ?

**A:** Yes, if  $f$  is nonsmooth at  $x$

**Q:** Can a function  $f$  have no subgradient at a point  $x$ ?

**A:** Yes, if  $x$  does not lie on convex hull of  $f$

## Subgradients and convexity

- ▶  $g$  is a subgradient of  $f$  at  $x$  iff  $(g, -1)$  supports **epi**  $f$  at  $(x, f(x))$
- ▶  $g$  is a subgradient iff  $f(x) + g^T(y - x)$  is a global (affine) underestimator of  $f$
- ▶ if  $f$  is convex and differentiable,  $\nabla f(x)$  is a subgradient of  $f$  at  $x$

subgradients come up in several contexts:

- ▶ algorithms for nondifferentiable convex optimization
- ▶ convex analysis, e.g., optimality conditions, duality for nondifferentiable problems

(if  $f(y) \leq f(x) + g^T(y - x)$  for all  $y$ , then  $g$  is a **supergradient**)

## Subdifferential

set of all subgradients of  $f$  at  $x$  is called the **subdifferential** of  $f$  at  $x$ , denoted  $\partial f(x)$

$$\partial f(x) = \{g : f(y) \geq f(x) + g^T(y - x) \quad \forall y\}$$

for any  $f$ ,

- ▶  $\partial f(x)$  is a closed convex set (can be empty)
- ▶  $\partial f(x) = \emptyset$  if  $f(x) = \infty$

proof: use the definition

## Subdifferential

set of all subgradients of  $f$  at  $x$  is called the **subdifferential** of  $f$  at  $x$ , denoted  $\partial f(x)$

$$\partial f(x) = \{g : f(y) \geq f(x) + g^T(y - x) \quad \forall y\}$$

for any  $f$ ,

- ▶  $\partial f(x)$  is a closed convex set (can be empty)
- ▶  $\partial f(x) = \emptyset$  if  $f(x) = \infty$

proof: use the definition

if  $f$  is convex,

- ▶  $\partial f(x)$  is nonempty, for  $x \in \text{relint dom } f$
- ▶  $\partial f(x) = \{\nabla f(x)\}$ , if  $f$  is differentiable at  $x$
- ▶ if  $\partial f(x) = \{g\}$ , then  $f$  is differentiable at  $x$  and  $g = \nabla f(x)$

## Compute subgradient via definition

$g \in \partial f(x)$  iff

$$f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(x) = |x|$  for  $x \in \mathbf{R}$ . suppose  $s \in \text{sign}(x)$ , where

$$\text{sign}(x) = \begin{cases} \{1\} & x > 0 \\ [-1, 1] & x = 0 \\ -\{1\} & x < 0. \end{cases}$$

then

$$f(y) = \max(y, -y) \geq sy = s(x + y - x) = |x| + s(y - x)$$

## Compute subgradient via definition

$g \in \partial f(x)$  iff

$$f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(x) = |x|$  for  $x \in \mathbf{R}$ . suppose  $s \in \text{sign}(x)$ , where

$$\text{sign}(x) = \begin{cases} \{1\} & x > 0 \\ [-1, 1] & x = 0 \\ -\{1\} & x < 0. \end{cases}$$

then

$$f(y) = \max(y, -y) \geq sy = s(x + y - x) = |x| + s(y - x)$$

so  $\text{sign}(x) \subseteq \partial f(x)$  (in fact, holds with equality)

## Compute subgradient via definition

$g \in \partial f(x)$  iff

$$f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(x) = |x|$  for  $x \in \mathbf{R}$ . suppose  $s \in \text{sign}(x)$ , where

$$\text{sign}(x) = \begin{cases} \{1\} & x > 0 \\ [-1, 1] & x = 0 \\ -\{1\} & x < 0. \end{cases}$$

then

$$f(y) = \max(y, -y) \geq sy = s(x + y - x) = |x| + s(y - x)$$

so  $\text{sign}(x) \subseteq \partial f(x)$  (in fact, holds with equality)

picture



## Compute subgradient via definition

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(x) = \max_i a_i^T x + b_i$ .

## Compute subgradient via definition

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(x) = \max_i a_i^T x + b_i$ . then for any  $i$ ,

$$\begin{aligned} f(y) &= \max_i a_i^T y + b_i \\ &\geq a_i^T y + b_i \\ &= a_i^T (x + y - x) + b_i \\ &= a_i^T x + b_i + a_i^T (y - x) \\ &= f(x) + a_i^T (y - x), \end{aligned}$$

where the last line holds for  $i \in \operatorname{argmax}_j a_j^T x + b_j$ . so

- ▶  $a_i \in \partial f(x)$  for each  $i \in \operatorname{argmax}_j a_j^T x + b_j$
- ▶  $\partial f(x)$  is convex, so

$$\operatorname{Co}\{a_i : i \in \operatorname{argmax}_j a_j^T x + b_j\} \subseteq \partial f(x)$$

## Compute subgradient via definition

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(X) = \lambda_{\max}(X)$ .

## Compute subgradient via definition

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**example.** let  $f(X) = \lambda_{\max}(X)$ . then

$$\begin{aligned} f(Y) &= \sup_{\|v\| \leq 1} v^T Y v \\ &= \sup_{\|v\| \leq 1} v^T (X + Y - X) v, \quad \|v\| \leq 1 \\ &= \sup_{\|v\| \leq 1} \left( v^T X v + v^T (Y - X) v \right), \quad \|v\| \leq 1 \\ &= v^T X v + \text{tr}(v v^T (Y - X)), \quad v \in \underset{\|v\| \leq 1}{\text{argmax}} v^T X v \\ &= \lambda_{\max}(X) + \text{tr}(v v^T (Y - X)), \quad v \in \underset{\|v\| \leq 1}{\text{argmax}} v^T X v \end{aligned}$$

- ▶  $v v^T \in \partial f(X)$  for each  $v \in \underset{\|v\| \leq 1}{\text{argmax}} v^T X v$
- ▶  $\partial f(x)$  is convex, so

$$\text{Co}\{v v^T : v \in \underset{\|v\| \leq 1}{\text{argmax}} v^T X v\} \subseteq \partial f(x)$$

# Outline

Subgradients

**Subgradient properties**

Subgradient method

Proximal operators

Proximal gradient method

Relations

Fixed points

Averaged operators

## Properties of subgradients

subgradient inequality:

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \mathbf{dom}(f)$$

for convex  $f$ , we'll show

- ▶ subgradients are monotone: for any  $x, y \in \mathbf{dom} f$ ,  $g_y \in \partial f(y)$ , and  $g_x \in \partial f(x)$ ,

$$(g_y - g_x)^T(y - x) \geq 0$$

- ▶  $\partial f(x)$  is continuous: if  $f$  is (lower semi-)continuous,  $x^{(k)} \rightarrow x$ ,  $g^{(k)} \rightarrow g$ , and  $g^{(k)} \in \partial f(x^{(k)})$  for each  $k$ , then  $g \in \partial f(x)$
- ▶  $\partial f(x) = \operatorname{argmax} g^T x - f(x)$

these will help us compute subgradients

## Subgradients are monotone

**fact.** for any  $x, y \in \text{dom } f$ ,  $g_y \in \partial f(y)$ , and  $g_x \in \partial f(x)$ ,

$$(g_y - g_x)^T (y - x) \geq 0$$

**proof.** same as for differentiable case:

$$f(y) \geq f(x) + g_x^T (y - x) \quad f(x) \geq f(y) + g_y^T (x - y)$$

add these to get

$$(g_y - g_x)^T (y - x) \geq 0$$

## Subgradients are preserved under limits

subgradient inequality:

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**fact.** if  $f$  is (lower semi-)continuous,  $x^{(k)} \rightarrow x$ ,  $g^{(k)} \rightarrow g$ , and  $g^{(k)} \in \partial f(x^{(k)})$  for each  $k$ , then  $g \in \partial f(x)$

**proof.**



## Subgradients are preserved under limits

subgradient inequality:

$$g \in \partial f(x) \iff f(y) \geq f(x) + g^T(y - x) \quad \forall y \in \text{dom}(f)$$

**fact.** if  $f$  is (lower semi-)continuous,  $x^{(k)} \rightarrow x$ ,  $g^{(k)} \rightarrow g$ , and  $g^{(k)} \in \partial f(x^{(k)})$  for each  $k$ , then  $g \in \partial f(x)$

**proof.** For each  $k$  and for every  $y$ ,

$$\begin{aligned} f(y) &\geq f(x^{(k)}) + (g^{(k)})^T(y - x^{(k)}) \\ \lim_{k \rightarrow \infty} f(y) &\geq \lim_{k \rightarrow \infty} f(x^{(k)}) + (g^{(k)})^T(y - x^{(k)}) \\ f(y) &\geq f(x) + g^T(y - x) \end{aligned}$$

**moral.** To find a subgradient  $g \in \partial f(x)$ , find points  $x^{(k)} \rightarrow x$  where  $f$  is differentiable, and let  $g = \lim_{k \rightarrow \infty} \nabla f(x^{(k)})$ .

## Subgradients are preserved under limits: example

consider  $f(x) = |x|$ . we know

$$\partial f(x) = \begin{cases} \{-1\} & x < 0 \\ ? & x = 0 \\ \{1\} & x > 0 \end{cases}$$

so

- ▶  $\lim_{x \rightarrow 0^+} \nabla(x) = 1$
- ▶  $\lim_{x \rightarrow 0^-} \nabla(x) = -1$

hence

## Subgradients are preserved under limits: example

consider  $f(x) = |x|$ . we know

$$\partial f(x) = \begin{cases} \{-1\} & x < 0 \\ ? & x = 0 \\ \{1\} & x > 0 \end{cases}$$

so

- ▶  $\lim_{x \rightarrow 0^+} \nabla(x) = 1$
- ▶  $\lim_{x \rightarrow 0^-} \nabla(x) = -1$

hence

- ▶  $-1 \in \partial f(0)$  and  $1 \in \partial f(0)$
- ▶  $\partial f(0)$  is convex, so  $[-1, 1] \subseteq \partial f(0)$
- ▶ and  $\partial f(0)$  is monotone, so  $[-1, 1] = \partial f(0)$

## Convex functions can't be very non-differentiable

### Theorem

*Rockafellar, Convex Analysis, Thm 25.5 a convex function  $f$  is differentiable almost everywhere on the interior of its domain.*

## Convex functions can't be very non-differentiable

### Theorem

*Rockafellar, Convex Analysis, Thm 25.5 a convex function  $f$  is differentiable almost everywhere on the interior of its domain.*

**corollary:** pick  $x \in \text{dom } f$  uniformly at random. then  $f$  is differentiable at  $x$  w/prob 1.

## Convex functions can't be very non-differentiable

### Theorem

*Rockafellar, Convex Analysis, Thm 25.5 a convex function  $f$  is differentiable almost everywhere on the interior of its domain.*

**corollary:** pick  $x \in \text{dom } f$  uniformly at random. then  $f$  is differentiable at  $x$  w/prob 1.

**corollary:** For a convex function  $f$  and any  $x$ , there is a sequence of points  $x^{(k)} \rightarrow x$  where  $f$  is differentiable.

## Subgradients and fenchel conjugates

**fact.**  $g \in \partial f(x) \iff f^*(g) + f(x) = g^T x$

(recall the conjugate function  $f^*(g) = \sup_x g^T x - f(x)$ .)

## Subgradients and fenchel conjugates

**proof.** if  $f^*(g) + f(x) = g^T x$ ,

$$\begin{aligned} f^*(g) &= \sup_y g^T y - f(y) \\ &\geq g^T y - f(y) \quad \forall y \\ f(y) &\geq g^T y - f^*(g) \quad \forall y \\ &= g^T y - g^T x + f(x) \quad \forall y \\ &= g^T (y - x) + f(x) \quad \forall y \end{aligned}$$

so  $g \in \partial f(x)$ . conversely, if  $g \in \partial f(x)$ ,



## Subgradients and fenchel conjugates

**proof.** if  $f^*(g) + f(x) = g^T x$ ,

$$\begin{aligned}f^*(g) &= \sup_y g^T y - f(y) \\&\geq g^T y - f(y) \quad \forall y \\f(y) &\geq g^T y - f^*(g) \quad \forall y \\&= g^T y - g^T x + f(x) \quad \forall y \\&= g^T (y - x) + f(x) \quad \forall y\end{aligned}$$

so  $g \in \partial f(x)$ . conversely, if  $g \in \partial f(x)$ ,

$$\begin{aligned}f(y) &\geq g^T (y - x) + f(x) \\g^T x - f(x) &\geq g^T y - f(y) \\\sup_y g^T x - f(x) &\geq \sup_y g^T y - f(y) \\g^T x - f(x) &\geq f^*(g)\end{aligned}$$

## Subgradients and fenchel conjugates

### Conclusion.

$$\begin{aligned}g \in \partial f(x) &\iff f^*(g) + f(x) = g^T x \\ &\iff x \in \operatorname{argmax}_x g^T x - f(x)\end{aligned}$$

consider the same implications for the function  $f^*$ :

$$\begin{aligned}x \in \partial f^*(g) &\iff f(x) + f^*(g) = x^T g \\ &\iff g \in \operatorname{argmax}_g g^T x - f^*(g)\end{aligned}$$

so all these conditions are equivalent, and

$$g \in \partial f(x) \iff x \in \partial f^*(g)!$$

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(x) = \|x\|_1$ . compute

$$f^*(g) = \sup_x g^T x - \|x\|_1$$

=

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(x) = \|x\|_1$ . compute

$$\begin{aligned} f^*(g) &= \sup_x g^T x - \|x\|_1 \\ &= \begin{cases} 0 & \|g\|_\infty \leq 1 \\ \infty & \text{otherwise} \end{cases} \end{aligned}$$

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(x) = \|x\|_1$ . compute

$$\begin{aligned} f^*(g) &= \sup_x g^T x - \|x\|_1 \\ &= \begin{cases} 0 & \|g\|_\infty \leq 1 \\ \infty & \text{otherwise} \end{cases} \end{aligned}$$

given  $x$ ,

$$\begin{aligned} \partial f(x) &= \operatorname{argmax}_g g^T x - f^*(g) \\ &= \operatorname{argmax}_{\|g\|_\infty \leq 1} g^T x \\ &= \mathbf{sign}(x) \end{aligned}$$

where **sign** is computed elementwise.

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(X) = \|X\|_*$ . compute

$$f^*(G) = \sup_X \operatorname{tr}(G, X) - \|X\|_*$$

=

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(X) = \|X\|_*$ . compute

$$\begin{aligned} f^*(G) &= \sup_X \operatorname{tr}(G, X) - \|X\|_* \\ &= \begin{cases} 0 & \|G\| \leq 1 \\ \infty & \text{otherwise} \end{cases} \end{aligned}$$

where  $\|G\| = \sigma_1(G)$  is the operator norm of  $G$ .

## Compute subgradient via fenchel conjugate

$$\partial f(x) = \operatorname{argmax}_g g^T x - f^*(g)$$

**example.** let  $f(X) = \|X\|_*$ . compute

$$\begin{aligned} f^*(G) &= \sup_X \operatorname{tr}(G, X) - \|X\|_* \\ &= \begin{cases} 0 & \|G\| \leq 1 \\ \infty & \text{otherwise} \end{cases} \end{aligned}$$

where  $\|G\| = \sigma_1(G)$  is the operator norm of  $G$ .

given  $X = U \mathbf{diag}(\sigma) V^T$ ,

$$\begin{aligned} \partial f(x) &= \operatorname{argmax}_G \operatorname{tr}(G, X) - f^*(G) \\ &= \operatorname{argmax}_{\|G\| \leq 1} \operatorname{tr}(G, X) \\ &= U \mathbf{diag}(\mathbf{sign}(\sigma)) V^T \end{aligned}$$

where **sign** is computed elementwise.



# Outline

Subgradients

Subgradient properties

**Subgradient method**

Proximal operators

Proximal gradient method

Relations

Fixed points

Averaged operators

## Subgradient method

the **subgradient method** is a simple algorithm to minimize nondifferentiable convex function  $f$

$$x^{(k+1)} = x^{(k)} - \alpha_k g^{(k)}$$

- ▶  $x^{(k)}$  is the  $k$ th iterate
- ▶  $g^{(k)}$  is **any** subgradient of  $f$  at  $x^{(k)}$
- ▶  $\alpha_k > 0$  is the  $k$ th step size

## Subgradient method

the **subgradient method** is a simple algorithm to minimize nondifferentiable convex function  $f$

$$x^{(k+1)} = x^{(k)} - \alpha_k g^{(k)}$$

- ▶  $x^{(k)}$  is the  $k$ th iterate
- ▶  $g^{(k)}$  is **any** subgradient of  $f$  at  $x^{(k)}$
- ▶  $\alpha_k > 0$  is the  $k$ th step size

**warning:** subgradient method is **not** a descent method.

## Subgradient method

the **subgradient method** is a simple algorithm to minimize nondifferentiable convex function  $f$

$$x^{(k+1)} = x^{(k)} - \alpha_k g^{(k)}$$

- ▶  $x^{(k)}$  is the  $k$ th iterate
- ▶  $g^{(k)}$  is **any** subgradient of  $f$  at  $x^{(k)}$
- ▶  $\alpha_k > 0$  is the  $k$ th step size

**warning:** subgradient method is **not** a descent method.  
instead, keep track of best point so far

$$f_{\text{best}}^{(k)} = \min_{i=1,\dots,k} f(x^{(i)})$$

## How to avoid slow convergence

don't use subgradient method for very high accuracy!

instead,

- ▶ for high accuracy: rewrite problem as LP or SDP; use IPM
- ▶ for medium accuracy:
  - ▶ regularize your objective (so it's strongly convex)

$$\tilde{f}(x) = f(x) + \alpha \|x - x^0\|^2$$

- ▶ smooth your objective (so it's smooth)

$$\tilde{f}(x) = \mathbb{E}_{y: \|y-x\| \leq \delta} f(y)$$

- ▶ infimal convolution (so it's smooth and strongly convex):

$$\tilde{f}(x) = \inf_y f(y) + \frac{\rho}{2} \|y - x\|^2$$

- ▶ more on these later...
  - ▶ for low accuracy: use a constant step size; terminate when you stop improving much or get bored

# Outline

Subgradients

Subgradient properties

Subgradient method

**Proximal operators**

Proximal gradient method

Relations

Fixed points

Averaged operators

## Proximal operator

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} \left( f(z) + \frac{1}{2} \|z - x\|_2^2 \right)$$

## Proximal operator

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} \left( f(z) + \frac{1}{2} \|z - x\|_2^2 \right)$$

►  $\mathbf{prox}_f : \mathbf{R}^d \rightarrow \mathbf{R}^d$



## Proximal operator

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} \left( f(z) + \frac{1}{2} \|z - x\|_2^2 \right)$$

- ▶  $\mathbf{prox}_f : \mathbf{R}^d \rightarrow \mathbf{R}^d$
- ▶ **generalized projection:** if  $\mathbf{1}_C$  is the indicator of set  $C$ ,

$$\mathbf{prox}_{\mathbf{1}_C}(w) = \Pi_C(w)$$

## Proximal operator

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $\mathbf{prox}_f : \mathbf{R}^d \rightarrow \mathbf{R}^d$
- ▶ **generalized projection:** if  $\mathbf{1}_C$  is the indicator of set  $C$ ,

$$\mathbf{prox}_{\mathbf{1}_C}(w) = \Pi_C(w)$$

- ▶ **implicit gradient step:** if  $z = \mathbf{prox}_f(x)$

$$\begin{aligned} \partial f(z) + z - x &= 0 \\ z &= x - \partial f(z) \end{aligned}$$

## Maps from functions to functions

no consistent notation for map from functions to functions.

for a function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$ ,

- ▶ **prox** maps  $f$  to a new function  $\mathbf{prox}_f : \mathbf{R}^d \rightarrow \mathbf{R}^d$ 
  - ▶  $\mathbf{prox}_f(x)$  evaluates this function at the point  $x$
- ▶  $\nabla$  maps  $f$  to a new function  $\nabla f : \mathbf{R}^d \rightarrow \mathbf{R}^d$ 
  - ▶  $\nabla f(x)$  evaluates this function at the point  $x$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

►  $f(x) = 0$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$



## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$  (separable)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$  (separable)
- ▶  $f(x) = \|x\|_1$

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$  (separable)
- ▶  $f(x) = \|x\|_1$  (soft-thresholding on each index)

## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$  (separable)
- ▶  $f(x) = \|x\|_1$  (soft-thresholding on each index)
- ▶  $f(X) = \|X\|_*$



## Let's evaluate some proximal operators!

define the **proximal operator** of the function  $f : \mathbf{R}^d \rightarrow \mathbf{R}$

$$\mathbf{prox}_f(x) = \underset{z}{\operatorname{argmin}} (f(z) + \frac{1}{2} \|z - x\|_2^2)$$

- ▶  $f(x) = 0$  (identity)
- ▶  $f(x) = x^2$  (shrinkage)
- ▶  $f(x) = |x|$  (soft-thresholding)
- ▶  $f(x) = \mathbf{1}(x \geq 0)$  (projection)
- ▶  $f(x) = \sum_{i=1}^d f_i(x_i)$  (separable)
- ▶  $f(x) = \|x\|_1$  (soft-thresholding on each index)
- ▶  $f(X) = \|X\|_*$  (soft-thresholding on singular values)

## Proxable functions

we say a function  $f$  is **proxable** if it's easy to evaluate  $\mathbf{prox}_f(x)$

all examples from previous slide are proxable

# Outline

Subgradients

Subgradient properties

Subgradient method

Proximal operators

**Proximal gradient method**

Relations

Fixed points

Averaged operators

## Proximal gradient method

suppose  $f$  is smooth,  $g$  is non-smooth. solve

$$\text{minimize } f(x) + g(x)$$

using proximal operators together with gradient steps?

## Proximal gradient method

suppose  $f$  is smooth,  $g$  is non-smooth. solve

$$\text{minimize } f(x) + g(x)$$

using proximal operators together with gradient steps? idea:

$$x^+ = \mathbf{prox}_{tg}(x - t\nabla f(x))$$

- ▶ the proximal operator gives a **fast method** to step towards the minimum of  $g$
- ▶ gradient method works well to step towards minimum of  $f$

## Proximal gradient: examples

- ▶ projected gradient  $g(x) = \mathbf{1}(\Omega)(x)$
- ▶ nonnegative least squares:  $f(x) = \frac{1}{2}\|Ax - b\|_2^2$ ,  
 $g(x) = \lambda\|x\|_1$
- ▶ lasso:  $f(x) = \frac{1}{2}\|Ax - b\|_2^2$ ,  $g(x) = \lambda\|x\|_1$
- ▶ ...

# Outline

Subgradients

Subgradient properties

Subgradient method

Proximal operators

Proximal gradient method

**Relations**

Fixed points

Averaged operators

# Functions

in much of what follows, we'll need to assume functions are

- ▶ closed: **epi**( $f$ ) is a closed set
- ▶ convex:  $f$  is convex
- ▶ proper: **dom**  $f$  is non-empty

which we abbreviate as CCP



## Relations

$(x, \partial f(x))$  and  $(x, \mathbf{prox}_f(x))$  define **relations** on  $\mathbf{R}^n$

- ▶ a **relation**  $R$  on  $\mathbf{R}^n$  is a subset of  $\mathbf{R}^n \times \mathbf{R}^n$
- ▶  $\mathbf{dom} R = \{x : (x, y) \in R\}$
- ▶ let  $R(x) = \{y : (x, y) \in R\}$
- ▶ if  $R(x)$  is always empty or a singleton, we say  $R$  is a function
- ▶ any function  $f : \mathbf{R}^n \rightarrow \mathbf{R}^n$  defines a relation  $\{(x, f(x)) : x \in \mathbf{dom} f\}$

## Relations: examples

- ▶ empty relation:  $\emptyset$
- ▶ full relation:  $\mathbf{R}^n \times \mathbf{R}^n$
- ▶ identity:  $\{(x, x) : x \in \mathbf{R}^n\}$
- ▶ zero:  $\{(x, 0) : x \in \mathbf{R}^n\}$
- ▶ subdifferential:  $\{(x, g) : x \in \mathbf{dom} f, g \in \partial f(x)\}$

## Operations on relations

if  $R$  and  $S$  are relations, define

- ▶ composition:  $RS = \{(x, z) : (x, y) \in R, (y, z) \in S\}$
- ▶ addition:  $R + S = \{(x, y + z) : (x, y) \in R, (x, z) \in S\}$
- ▶ inverses:  $R^{-1} = \{(y, x) : (x, y) \in R\}$

use inequality on sets to mean the inequality holds for any element in the set, e.g.,

$$f(y) \geq f(x) + \partial f^T(y - x)$$

## Example: fenchel conjugates and the subdifferential

if  $f$  is CPP,  $(f^*)^* = f^{**} = f$ , so

$$\begin{aligned}(u, v) \in (\partial f)^{-1} &\iff (v, u) \in \partial f \\&\iff u \in \partial f(v) \\&\iff 0 \in \partial f(v) - u \\&\iff v \in \operatorname{argmin}_x (f(x) - u^T x) \\&\iff v \in \operatorname{argmax}_x (u^T x - f(x)) \\&\iff f(v) + f^*(u) = u^T v \\&\iff u \in \operatorname{argmax}_y (y^T v - f^*(y)) \\&\iff 0 \in v - \partial f^*(u) \\&\iff (u, v) \in \partial f^*\end{aligned}$$

this shows  $\partial f^* = \partial f^{-1}$

# Outline

Subgradients

Subgradient properties

Subgradient method

Proximal operators

Proximal gradient method

Relations

**Fixed points**

Averaged operators

## Zeros of a relation

- ▶  $x$  is a **zero** of  $R$  if  $0 \in R(x)$
- ▶ the **zero set** of  $R$  is  $R^{-1}(0) = \{x : (x, 0) \in R\}$

## Zeros of a relation

- ▶  $x$  is a **zero** of  $R$  if  $0 \in R(x)$
- ▶ the **zero set** of  $R$  is  $R^{-1}(0) = \{x : (x, 0) \in R\}$

$x$  is a zero of  $\partial f$  iff  $x$  solves minimize  $f(x)$

## Lipschitz operators

relation  $F$  has Lipschitz constant  $L$  if for all  $(x, u) \in F$  and  $(y, v) \in F$ ,

$$\|u - v\| \leq L\|x - y\|$$

**fact:** if  $F$  is Lipschitz, then  $F$  is a function.

**proof:**



## Lipschitz operators

relation  $F$  has Lipschitz constant  $L$  if for all  $(x, u) \in F$  and  $(y, v) \in F$ ,

$$\|u - v\| \leq L\|x - y\|$$

**fact:** if  $F$  is Lipschitz, then  $F$  is a function.

**proof:** if  $(x, u) \in F$  and  $(x, v) \in F$ ,

$$\|u - v\| \leq L\|x - x\| = 0$$

- ▶ the relation  $F$  is **nonexpansive** if  $L \leq 1$
- ▶ the relation  $F$  is **contractive** if  $L < 1$

## Gradient update is contractive for SSC functions

suppose  $f$  is  $\alpha$ -strongly convex and  $\beta$ -smooth. the relation

$$I - t\nabla f = \{(x, x - t\nabla f(x)) : x \in \mathbf{dom} f\}$$

is Lipschitz with parameter  $L = \max\{|1 - t\alpha|, |1 - t\beta|\}$ .

## Gradient update is contractive for SSC functions

suppose  $f$  is  $\alpha$ -strongly convex and  $\beta$ -smooth. the relation

$$I - t\nabla f = \{(x, x - t\nabla f(x)) : x \in \mathbf{dom} f\}$$

is Lipschitz with parameter  $L = \max\{|1 - t\alpha|, |1 - t\beta|\}$ .

**corollary:** if  $t = \frac{2}{\alpha + \beta}$ ,  $L = \frac{\kappa - 1}{\kappa + 1}$

## Gradient update is contractive for SSC functions

suppose  $f$  is  $\alpha$ -strongly convex and  $\beta$ -smooth. the relation

$$I - t\nabla f = \{(x, x - t\nabla f(x)) : x \in \mathbf{dom} f\}$$

is Lipschitz with parameter  $L = \max\{|1 - t\alpha|, |1 - t\beta|\}$ .

**corollary:** if  $t = \frac{2}{\alpha + \beta}$ ,  $L = \frac{\kappa - 1}{\kappa + 1}$

**hint:** use the fundamental theorem of calculus

$$(I - t\nabla f)(x) - (I - t\nabla f)(y) = \int_0^1 (I - t\nabla^2 f(\theta x + (1 - \theta)y))(x - y) dt$$

and Jensen's inequality

$$\left\| \int_0^1 v(t) dt \right\| \leq \int_0^1 \|v(t)\| dt$$

source: Ryu and Yin (2022)

## Gradient update is contractive for SSC functions

suppose  $f$  is  $\alpha$ -strongly convex and  $\beta$ -smooth. the relation

$$I - t\nabla f = \{(x, x - t\nabla f(x)) : x \in \mathbf{dom} f\}$$

is Lipschitz with parameter  $L = \max\{|1 - t\alpha|, |1 - t\beta|\}$ .

## Gradient update is contractive for SSC functions

suppose  $f$  is  $\alpha$ -strongly convex and  $\beta$ -smooth. the relation

$$I - t\nabla f = \{(x, x - t\nabla f(x)) : x \in \mathbf{dom} f\}$$

is Lipschitz with parameter  $L = \max\{|1 - t\alpha|, |1 - t\beta|\}$ .

**proof:**

$$\begin{aligned} & \| (I - t\nabla f)(x) - (I - t\nabla f)(y) \| \\ &= \left\| \int_0^1 (I - t\nabla^2 f(\theta x + (1 - \theta)y))(x - y) dt \right\| \\ &\leq \int_0^1 \| (I - t\nabla^2 f(\theta x + (1 - \theta)y))(x - y) \| dt \\ &\leq \int_0^1 \max(|1 - t\alpha|, |1 - t\beta|) dt \|x - y\| \\ &= \max(|1 - t\alpha|, |1 - t\beta|) \|x - y\| \end{aligned}$$

## Proximal map is nonexpansive

the proximal map of any convex function  $f$  is nonexpansive:

$$\|\mathbf{prox}_f(y) - \mathbf{prox}_f(x)\| \leq \|y - x\|$$

## Proximal map is nonexpansive

the proximal map of any convex function  $f$  is nonexpansive:

$$\|\mathbf{prox}_f(y) - \mathbf{prox}_f(x)\| \leq \|y - x\|$$

**proof:** let  $u = \mathbf{prox}_f(x)$  and  $v = \mathbf{prox}_f(y)$ , so

$$x - u \in \partial f(u), \quad y - v \in \partial f(v)$$

then by the subgradient inequality,

$$f(v) \geq f(u) + \langle x - u, v - u \rangle \quad \text{and} \quad f(u) \geq f(v) + \langle y - v, u - v \rangle$$

add these to show

$$\begin{aligned} 0 &\geq \langle y - x + u - v, u - v \rangle \\ \langle x - y, u - v \rangle &\geq \|u - v\|^2 \\ \|x - y\| &\geq \|u - v\| \end{aligned}$$

- ▶ second line shows  $\mathbf{prox}_f$  is firmly nonexpansive
- ▶ third line uses Cauchy-Schwarz to show it is nonexpansive



## Proximal map is contractive for SC functions

the proximal map of an  $\alpha$ -SC function  $f$  is  $\frac{1}{1+2\alpha}$ -contractive:

$$\|\mathbf{prox}_f(y) - \mathbf{prox}_f(x)\| \leq \frac{1}{1+2\alpha} \|y - x\|$$

## Proximal map is contractive for SC functions

the proximal map of an  $\alpha$ -SC function  $f$  is  $\frac{1}{1+2\alpha}$ -contractive:

$$\|\mathbf{prox}_f(y) - \mathbf{prox}_f(x)\| \leq \frac{1}{1+2\alpha} \|y - x\|$$

**proof:** let  $u = \mathbf{prox}_f(x)$  and  $v = \mathbf{prox}_f(y)$ , so

$$x - u \in \partial f(u), \quad y - v \in \partial f(v)$$

by strong convexity

$$\begin{aligned} f(v) &\geq f(u) + \langle x - u, v - u \rangle + \alpha \|v - u\|^2 \\ f(u) &\geq f(v) + \langle y - v, u - v \rangle + \alpha \|u - v\|^2 \end{aligned}$$

add these to show

$$\begin{aligned} 0 &\geq \langle y - x + u - v, u - v \rangle + 2\alpha \|u - v\|^2 \\ \langle x - y, u - v \rangle &\geq (1 + 2\alpha) \|u - v\|^2 \\ \frac{1}{1 + 2\alpha} \|x - y\| &\geq \|u - v\| \end{aligned}$$

## Fixed points

$x$  is a **fixed point** of  $F$  if  $x = F(x)$

examples:

- ▶  $F(x) = x$ : every point is a fixed point
- ▶  $F(x) = 0$ : only 0 is a fixed point

## Fixed points

$x$  is a **fixed point** of  $F$  if  $x = F(x)$

examples:

- ▶  $F(x) = x$ : every point is a fixed point
- ▶  $F(x) = 0$ : only 0 is a fixed point
- ▶ a contractive operator on  $\mathbf{R}^n$  can have at most one FP

## Fixed points

$x$  is a **fixed point** of  $F$  if  $x = F(x)$

examples:

- ▶  $F(x) = x$ : every point is a fixed point
- ▶  $F(x) = 0$ : only 0 is a fixed point
- ▶ a contractive operator on  $\mathbf{R}^n$  can have at most one FP

**proof:** if  $x$  and  $y$  are FPs,

$$\|x - y\| = \|F(x) - F(y)\| < \|x - y\| \text{ contradiction}$$

## Fixed points

$x$  is a **fixed point** of  $F$  if  $x = F(x)$

examples:

- ▶  $F(x) = x$ : every point is a fixed point
- ▶  $F(x) = 0$ : only 0 is a fixed point
- ▶ a contractive operator on  $\mathbf{R}^n$  can have at most one FP  
**proof:** if  $x$  and  $y$  are FPs,  
 $\|x - y\| = \|F(x) - F(y)\| < \|x - y\|$  contradiction
- ▶ a nonexpansive operator  $F$  need not have a fixed point

## Fixed points

$x$  is a **fixed point** of  $F$  if  $x = F(x)$

examples:

- ▶  $F(x) = x$ : every point is a fixed point
- ▶  $F(x) = 0$ : only 0 is a fixed point
- ▶ a contractive operator on  $\mathbf{R}^n$  can have at most one FP  
**proof:** if  $x$  and  $y$  are FPs,  
 $\|x - y\| = \|F(x) - F(y)\| < \|x - y\|$  contradiction
- ▶ a nonexpansive operator  $F$  need not have a fixed point  
**proof:** translation

## Fixed point iteration

to find a fixed point of  $F$ , try the fixed point iteration

$$x^{(k+1)} = F(x^{(k)})$$



## Fixed point iteration

to find a fixed point of  $F$ , try the fixed point iteration

$$x^{(k+1)} = F(x^{(k)})$$

**Q:** when does this converge?

## Fixed point iteration: contractive

**Banach fixed point theorem:** if  $F$  is a contraction, the iteration

$$x^{(k+1)} = F(x^{(k)})$$

converges to the unique fixed point of  $F$

properties: if  $L$  is the Lipschitz constant of  $F$ ,

- ▶ distance to fixed point decreases monotonically:

$$\|x^{(k+1)} - x^*\| = \|F(x^{(k)}) - F(x^*)\| \leq L\|x^{(k)} - x^*\|$$

(iteration is **Fejer-monotone**)

- ▶ linear convergence with rate  $L$

# Proof

**proof:**

## Proof

**proof:** if  $F$  has Lipschitz constant  $L < 1$ ,

► sequence  $x^{(k)}$  is Cauchy:

$$\begin{aligned}\|x^{(k+\ell)} - x^{(k)}\| &\leq \|x^{(k+\ell)} - x^{(k+\ell-1)}\| + \dots + \|x^{(k+1)} - x^{(k)}\| \\ &\leq (L^{\ell-1} + \dots + 1)\|x^{(k+1)} - x^{(k)}\| \\ &\leq \frac{1}{1-L}\|x^{(k+1)} - x^{(k)}\| \\ &\leq \frac{L^k}{1-L}\|x^{(1)} - x^{(0)}\|\end{aligned}$$

► so it converges to a point  $x^*$ . must be the (unique) FP!

► converges to  $x^*$  linearly with rate  $L$

$$\|x^{(k)} - x^*\| = \|F(x^{(k-1)}) - F(x^*)\| \leq L\|x^{(k-1)} - x^*\| \leq L^k\|x^{(0)} - x^*\|$$

# Outline

Subgradients

Subgradient properties

Subgradient method

Proximal operators

Proximal gradient method

Relations

Fixed points

Averaged operators

## Fixed point iteration: nonexpansive

if  $F$  is nonexpansive, the iteration

$$x^{(k+1)} = F(x^{(k)})$$

need not converge to a fixed point even if one exists.

**proof:**

## Fixed point iteration: nonexpansive

if  $F$  is nonexpansive, the iteration

$$x^{(k+1)} = F(x^{(k)})$$

need not converge to a fixed point even if one exists.

**proof:**

- ▶ let  $F$  rotate its argument by  $\theta$  degrees around the origin.
- ▶ then  $F$  is nonexpansive and has a fixed point at  $x^* = 0$ .
- ▶ but if  $\|x^{(0)}\| = r$ , then  $\|F(x^{(k)})\| = r$  for all  $k$ .

## Averaged operators

an operator  $F$  is **averaged** if

$$F = \theta G + (1 - \theta)I$$

for  $\theta \in (0, 1)$ ,  $G$  nonexpansive



## Averaged operators

an operator  $F$  is **averaged** if

$$F = \theta G + (1 - \theta)I$$

for  $\theta \in (0, 1)$ ,  $G$  nonexpansive

**fact:** if  $F$  is averaged, then  $x$  is FP of  $F \iff x$  is FP of  $G$

**proof:**

## Averaged operators

an operator  $F$  is **averaged** if

$$F = \theta G + (1 - \theta)I$$

for  $\theta \in (0, 1)$ ,  $G$  nonexpansive

**fact:** if  $F$  is averaged, then  $x$  is FP of  $F \iff x$  is FP of  $G$

**proof:**

$$\begin{aligned}x &= Fx = \theta Gx + (1 - \theta)x = \theta Gx + (1 - \theta)x \\ \theta x &= \theta Gx \\ x &= Gx\end{aligned}$$

$\implies$  if  $G$  is nonexpansive,  $F = \frac{1}{2}I + \frac{1}{2}G$  is averaged with same FPs

## Fixed point iteration: averaged

if  $F = \theta G + (1 - \theta)I$  is averaged ( $\theta \in (0, 1)$ ,  $G$  nonexpansive), the iteration

$$x^{(k+1)} = F(x^{(k)})$$

converges to a fixed point if one exists.

(also called the damped, averaged, or Mann-Krasnosel'skii iteration.)

properties: Ryu and Boyd (2016)

- ▶ distance to fixed point decreases monotonically (Fejer-monotone)
- ▶ sublinear convergence of fixed point residual

$$\|Gx^{(k)} - x^{(k)}\|^2 \leq \frac{1}{(k+1)\theta(1-\theta)} \|x^{(0)} - x^*\|^2$$

## Gradient descent operator is averaged

follows Ryu and Yin (2022)

**fact:** if  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $\beta$ -smooth, then  $I - \frac{2}{\beta} \nabla f$  is non-expansive

## Gradient descent operator is averaged

follows Ryu and Yin (2022)

**fact:** if  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $\beta$ -smooth, then  $I - \frac{2}{\beta}\nabla f$  is non-expansive

**proof:** since  $f$  is  $\beta$ -smooth,

$$\begin{aligned}\|(I - \frac{2}{\beta}\nabla f)(x) - (I - \frac{2}{\beta}\nabla f)(y)\|^2 &= \|x - y\|^2 - \frac{4}{\beta} \left( \langle x - y, \nabla f(x) - \nabla f(y) \rangle \right) \\ &\leq \|x - y\|^2\end{aligned}$$

## Gradient descent operator is averaged

follows Ryu and Yin (2022)

**fact:** if  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $\beta$ -smooth, then  $I - \frac{2}{\beta}\nabla f$  is non-expansive

**proof:** since  $f$  is  $\beta$ -smooth,

$$\begin{aligned}\|(I - \frac{2}{\beta}\nabla f)(x) - (I - \frac{2}{\beta}\nabla f)(y)\|^2 &= \|x - y\|^2 - \frac{4}{\beta} \left( \langle x - y, \nabla f(x) - \nabla f(y) \rangle \right) \\ &\leq \|x - y\|^2\end{aligned}$$

**corollary:** if  $f : \mathbf{R}^n \rightarrow \mathbf{R}$  is  $\beta$ -smooth, then  $I - t\nabla f$  is averaged for  $t \in (0, \frac{2}{\beta})$

since  $I - t\nabla f = (1 - \frac{t\beta}{2})I + \frac{t\beta}{2}(I - \frac{2}{\beta}\nabla f)$

## When does proximal gradient converge?

proximal gradient converges at rate  $O(1/k)$  when  $I - t\nabla f$  is averaged and  $\mathbf{prox}_{tg}$  is nonexpansive

- ▶ if  $f$  is  $\beta$ -smooth and step size  $t \in (0, \frac{2}{\beta})$
- ▶ and  $g$  is convex

proximal gradient converges linearly when, in addition,  $I - t\nabla f$  or  $\mathbf{prox}_{tg}$  is contractive

- ▶ if  $f$  is  $\beta$ -smooth and  $\alpha$ -strongly convex and  $\max(|1 - t\alpha|, |1 - t\beta|) < 1$
- ▶ or if  $g$  is strongly convex